

Family Medicine


Appendix A: Demographics, Site #1




Site 1 offers a 5-year dual FM/psychiatry combined residency program. PGY345 includes all senior FM residents and senior FM/psych combined residents.



75% of participants had 2 or fewer experiences with diagnosing AUD in the preceding 3 months.



75% of participants had 2 or fewer experiences with performing a brief intervention or motivational interviewing in the preceding 3 months.



95% of participants had 2 or fewer experiences with prescribing medication for AUD in the preceding 3 months.

Family Medicine

Appendix B: Survey

Family Medicine

Part I: Physician confidence assessment, modeled on validated self – efficacy scales

1) How confident are you in identifying 'unhealthy alcohol use' in outpatient clinical practice?

1 – not at all confident | 2 – slightly confident | 3 – somewhat confident | 4 – neutral | 5 –moderately confident | 6 – highly confident | 7 – completely confident.

2) How confident are you in screening for unhealthy alcohol use/alcohol use disorder using a validated screening tool?

1 – not at all confident | 2 – slightly confident | 3 – somewhat confident | 4 – neutral | 5 –moderately confident | 6 – highly confident | 7 – completely confident.

3) How confident are you in applying a brief (MI) intervention for unhealthy alcohol use/alcohol use disorder?

1 – not at all confident | 2 – slightly confident | 3 – somewhat confident | 4 – neutral | 5 –moderately confident | 6 – highly confident | 7 – completely confident.

4) How confident are you in providing a referral to treatment for unhealthy alcohol use/alcohol use disorder?

1 – not at all confident | 2 – slightly confident | 3 – somewhat confident | 4 – neutral | 5 –moderately confident | 6 – highly confident | 7 – completely confident.

5) How confident are you in determining appropriate treatments for unhealthy alcohol use/alcohol use disorder?

1 – not at all confident | 2 – slightly confident | 3 – somewhat confident | 4 – neutral | 5 –moderately confident | 6 – highly confident | 7 – completely confident.

Part II: Physician knowledge assessment of SBIRT and outpatient AUD medication management

6) Define SBIRT:

7) List 2 validated screening tools for unhealthy alcohol use / alcohol use disorder:

8) A 34-year-old male drinks 4 alcoholic beverages twice a week. Is this a positive screen for unhealthy alcohol use?

9) A 44-year-old female drinks 4 alcoholic beverages once a week. Is this a positive screen for unhealthy alcohol use?

10) Describe the 4 components of the 'OARS' motivational interviewing framework:

11) List 1 referral to treatment option for unhealthy alcohol use / AUD in the family health centers:

For questions 12-16, match one medication to each clinical scenario (each medication will only be chosen once).

12) A 54 yoM with HTN, HLD, severe AUD, and OUD on methadone presents to discuss medication management after having recently completed alcohol detoxification. Answer: _____

13) A 32 yoF with severe AUD and recent admission for acute alcohol – induced hepatitis with repeat AST/ALT 350/120 presents to discuss a medication to reduce heavy drinking days. Answer: _____

14) A 28 yoF with IDA, GAD, MDD, and mild AUD presents to discuss medication management after having intolerance to vivitrol. Answer: _____

15) A 35 yoM with HTN, TUD, severe AUD with compensated cirrhosis presents to discuss a medication to improve alcohol abstinence. Answer: _____

16) A 30 yoF with migraines with aura, obesity, and mild AUD presents to discuss a medication to reduce the number of heavy drinking days. Answer: _____

A) Naltrexone
B) Acamprosate
C) Gabapentin
D) Sertraline
E) Topiramate

Key: HTN – hypertension; HLD – hyperlipidemia; AUD – alcohol use disorder; OUD – opioid use disorder; IDA – iron deficiency anemia; GAD – generalized anxiety disorder; MDD – major depressive disorder; TUD – tobacco use disorder.